CIRSIUM MUTICUM Michx., var. subpinnatifidum (Britton), n. comb. Carduus muticus, var. subpinnatifidus Britton in Britton & Brown, Ill. Fl. iii. 489 (1898).

Cirsium Hillii (Canby), n. comb. Cnicus Hillii Canby, Gard. & For. iv. 101 (1891). Carduus Hillii Porter, Mem. Torr. Bot. Cl. v. 344 (1894).

Prenanthes altissima L., var. cinnamomea, n. var., pappo cinnamomeo.— Pappus cinnamon-color; otherwise like the typical form.— Indiana to Missouri and Louisiana. Type collected in rich woods, Monteer, Missouri, October 5, 1905 (Bush, no. 3534).

GRAY HERBARIUM.

The Annual Botanical Symposium.— The fifth annual symposium will be held at Georgetown, Delaware, July 1–8, 1908. Head-quarters will be Hotel Eagle, where rooms may be secured at \$1.50 a day. Trips will be made to Lewes, Rehoboth, Chesapeake Bay, Indian River, etc. Those desiring to attend should inform Joseph Crawford, 2824 Frankford Ave., Philadelphia, Pa.

Josselyn Botanical Society of Maine will be held at Manset, Mount Desert Island, August 4th to 8th, 1908. A reduced rate of \$2.00 per day for board and lodging has been granted by the Stanley House, and reduced railroad fares will be arranged. All who propose attending should communicate at once with O. W. Knight, Chairman of Committee on Arrangements, Bangor, Maine.

Lemna minor and Sparganium Eurycarpum in Rimouski County, Quebec.— For several summers Professor J. F. Collins and the writer have visited Bic and the adjacent towns of Rimouski County, Quebec, on account of the great development in that region of bold limestone headlands and ridges, upon which are found an extensive high-northern flora (see Rhodora, ix. 158–172). Bordering the harbor of Bic is a salt marsh, the brackish borders of which likewise support a pronouncedly northern flora: Catabrosa aquatica (L.) Beauv., Poa eminens J. S. Presl, Scirpus rufus (Huds.) Schrad., Carex norvegica Willd., C. glareosa Wahlenb., Rumex occidentalis Watson, etc. In